# Index of Authors


ACHARYA, A. S. *see* MANJULA, B. N. 337–345
ANSTEE, D. J. *see* TANNER, M. J. A. 157–161
ARONSSON, A.-C. & MANNERVIK, B. Characterization of glyoxalase I purified from pig erythrocytes by affinity chromatography 503–509
ASHTON, A. R. & POLYA, G. M. Adenosine 3′:5′-cyclic monophosphate in higher plants. Isolation and characterization of adenosine 3′:5′-cyclic monophosphate from *Kalanchoe* and *Agave* 27–32
ATKINS, G. L. *see* NIMMO, I. A. 107–110
AWASTHI, Y. C. *see* WIKTOROWICZ, J. E. 49–53

BANNISTER, J. V. *see* CASS, A. E G. 587–589
BANNISTER, W. H. *see* CASS, A. E. G. 587–589
BARTHOLOMEW, B., DODGSON, K. S., MATCHAM, G. W. J., SHAW, D. J. & WHITE, G. F. A novel mechanism of enzymic ester hydrolysis. Inversion of configuration and carbon–oxygen bond cleavage by secondary alkylsulphohydrolases from detergent-degrading microorganisms 575–580
BJØRKLID, E. & STORM, E. Purification and some properties of the protein component of tissue thromboplastin from human brain 89–96
BLACKWELL, L. F. *see* MACGIBBON, A. K. H. 455–462
BOCCU', E. *see* FONTANA, A. 539–545
BOLLACK, C. *see* HELWIG, J.-J. 127–134
BOWLEY, M., COOLING, J., BURDITT, S. L. & BRINDLEY, D. N. The effects of amphiphilic cationic drugs and inorganic cations on the activity of phosphatidate phosphohydrolase 447–454
BRAZEAU, P. *see* OGAWA, N. 269–277
BRINDLEY, D. N. *see* BOWLEY, M. 447–454
BRITTAIN, T. *see* GREENWOOD, C. 413–416; THOMSON, A. J. 327–336
BRUNDISH, D. E. & WADE, R. The synthesis of tritium-labelled human corticotropin of high specific radioactivity 169–171
BRUNORI, M. *see* GREENWOOD, C. 413–416
BUCKLEY, P. D. *see* MACGIBBON, A. K. H. 455–462
BURDITT, S. L. *see* BOWLEY, M. 447–454
BURSTEIN, Y. & SCHECHTER, I. Glutamine as a precursor to *N*-terminal pyrrolid-2-one-5-carboxylic acid in mouse immunoglobulin $\lambda$-type light chains. Amino acid-sequence variability at the *N*-terminal extra piece of $\lambda$-type light-chain precursors 347–354

CARRELL, R. W. *see* WINTERBOURN, C. C. 141–148
CASS, A. E. G., HILL, H. A. O., SMITH, B. E., BANNISTER, J. V. & BANNISTER, W. H. Carbon-2 proton exchange at histidine-41 in bovine erythrocyte superoxide dismutase 587–589
CHAPMAN, D. J. *see* HEANEY-KIERAS, J. 1–9
CHIPKO, B. R. *see* CORBETT, M. D. 263–267
CHOONG, Y. S., SHEPHERD, M. G. & SULLIVAN, P. A. Modification of lactate oxidase with diethyl pyrocarbonate. Evidence for an active-site histidine residue 385–393
CHOONG, Y. S. *see also* SULLIVAN, P. A. 375–383
CLARKE, F. M. *see* WALSH, T. P. 165–167

COGGINS, J. R. *see* LA NAUZE, J. M. 409–411
COLBY, J., STIRLING, D. I. & DALTON, H. The soluble methane mono-oxygenase of *Methylococcus capsulatus* (Bath). Its ability to oxygenate *n*-alkanes, *n*-alkenes, ethers and alicyclic, aromatic and heterocyclic compounds 395–402
COOLING, J. *see* BOWLEY, M. 447–454
CORBETT, M. D. & CHIPKO, B. R. *N*-Phenylglycolhydroxamate production by the action of transketolase on nitrosobenzene 263–267
CORNISH-BOWDEN, A. An automatic method for deriving steady-state rate equations 55–59
CORNISH-BOWDEN, A. *see also* STORER, A. C. 61–69; THORNELEY, R. N. F. 255–262
CRAMB, R. *see* STRANGE, R. C. 425–429
CREASER, E. H. *see* LINDSAY, J. A. 247–253
CROWDER, S. E. & RAGAN, C. I. Effects of proteolytic digestion by chymotrypsin on the structure and catalytic properties of reduced nicotinamide–adenine dinucleotide–ubiquinone oxidoreductase from bovine heart mitochondria 295–301
CUTFIELD, J. F. *see* SULLIVAN, P. A. 375–383

DADAY, A. *see* SMITH, G. D. 121–126
DALTON, H. *see* COLBY, J. 395–402
DIXON, H. B. F. *see* LA NAUZE, J. M. 409–411
DODGSON, K. S. *see* BARTHOLOMEW, B. 575–580
DØSKELAND, S. O. & UELAND, P. M. Binding proteins for adenosine 3′:5′-cyclic monophosphate in bovine adrenal cortex 561–573
DOWER, S. K., WAIN-HOBSON, S., GETTINS, P., GIVOL, D., JACKSON, W. R. C., PERKINS, S. J., SUNDERLAND, C. A., SUTTON, B. J., WRIGHT, C. E. & DWEK, R. A. The combining site of the dinitrophenyl-binding immunoglobulin A myeloma protein MOPC 315 207–223
DOWER, S. K. *see also* PERKINS, S. J. 223–225; WAIN-HOBSON, S. 227–235
DUGAL, B. & STRØMME, J. Purification and some properties of 1-aspartamido-$\beta$-*N*-acetylglucosamine amidohydrolase from human liver 497–502
DURBAN, E. *see* NOCHUMSON, S. 11–18
DWEK, R. A. *see* DOWER, S. K. 207–223; PERKINS, S. J. 223–225; SUTTON, B. J. 177–197; WAIN-HOBSON, S. 227–235; WILLAN, K. J. 199–206

FAROOQUI, A. A. *see* HELWIG, J.-J. 127–134
FIELDEN, E. M. *see* MCADAM, M. E. 71–79, 81–87
FONTANA, A., VITA, C., BOCCU', E. & VERONESE, F. M. A fluorimetric study of the role of calcium ions in the stability of thermolysin 539–545
FOX, R. A. *see* MCADAM, M. E. 71–79, 81–87
FRIESEN, H. G. *see* OGAWA, N. 269–277
FROST, R. G. *see* MILLER, A. L. 591–594

GERWIG, G. J. *see* HALL, R. L. 173–176
GETTINS, P. *see* DOWER, S. K. 207–223; PERKINS, S. J. 223–225; SUTTON, B. J. 177–197; WAIN-HOBSON, S. 227–235

GIGLI, I., VON ZABERN, I. & PORTER, R. R. The isolation and structure of C4, the fourth component of human complement 439–446
GILES, I. G., POAT, P. C. & MUNDAY, K. A. An investigation of the interactions of the allosteric modifiers of pyruvate kinase with the enzyme from *Carcinus maenas* hepatopancreas 97–105
GIVOL, D. *see* DOWER, S. K. 207–223; SUTTON, B. J. 177–197; WAIN-HOBSON, S. 227–235; WILLAN, K. J. 199–206
GRANT, D. A. W. *see* MAGEE, A. I. 583–585
GREENWOOD, C., BRITTAIN, T., BRUNORI, M. & WILSON, M. T. Chromous ion reduction of mammalian cytochrome oxidase and some of its derivatives 413–416
GREENWOOD, C. *see also* THOMSON, A. J. 327–336

HALL, R. L., WOOD, E. J., KAMERLING, J. P., GERWIG, G. J. & VLIEGENTHART, J. F. G. 3-*O*-Methyl sugars as constituents of glycoproteins. Identification of 3-*O*-methylgalactose and 3-*O*-methylmannose in pulmonate gastropod haemocyanins 173–176
HARPER, D. B. Microbial metabolism of aromatic nitriles. Enzymology of C–N cleavage by *Nocardia* sp. (*Rhodochrous* group) N.C.I.B. 11216 309–319
HAYES, J. D. *see* STRANGE, R. C. 425–429
HEANEY-KIERAS, J., RODÉN, L. & CHAPMAN, D. J. The covalent linkage of protein to carbohydrate in the extracellular protein–polysaccharide from the red alga *Porphyridium cruentum* 1–9
HELWIG, J.-J., FAROOQUI, A. A., BOLLACK, C. & MANDEL, P. Purification and some properties of arylsulphatases A and B from rabbit kidney cortex 127–134
HERMON-TAYLOR, J. *see* MAGEE, A. I. 583–585
HICKMAN, P. E. *see* WEIDEMANN, M. J. 525–537
HILL, H. A. O. *see* CASS, A. E. G. 587–589
HOVINGH, P. & LINKER, A. Specificity of flavobacterial glycuronidases acting on disaccharides derived from glycosaminoglycans 287–293
HURST, P. L., NIELSEN, J., SULLIVAN, P. A. & SHEPHERD, M. G. Purification and properties of a cellulase from *Aspergillus niger* 33–41

ISSINGER, O.-G. Purification and properties of a ribosomal casein kinase from rabbit reticulocytes 511–518

JACKSON, W. R. C. *see* DOWER, S. K. 207–223
JONES, D. S. *see* OLD, J. M. 367–373
JUDSON, P. A. *see* TANNER, M. J. A. 157–161
JULIANO, L. *see* NOUAILHETAS, V. L. A. 547–551

KAMERLING, J. P. *see* HALL, R. L. 173–176
KEENAN, R. W., RICE, N. & QUOCK, R. The high-performance liquid-chromatographic analysis of ficaprenol and dolichol 405–408
KIENER, P. A. & WALEY, S. G. Substrate-induced deactivation of penicillinases. Studies of $\beta$-lactamase I by hydrogen exchange 279–285
KIM, S. *see* NOCHUMSON, S. 11–18
KOLBUCH-BRADDON, M. E. *see* WEIDEMANN, M. J. 525–537
KONDO, M. *see* MOENS, L. 111–119
KURIHARA, T., NISHIZAWA, Y. & TAKAHASHI, Y. The use of non-aqueous chloroform/methanol extraction for the delipidation of brain with minimal loss of enzyme activities 135–140
KUROSKY, A. *see* WIKTOROWICZ, J. E. 49–53

LA NAUZE, J. M., COGGINS, J. R. & DIXON, H. B. F. Aldolase-like imine formation in the mechanism of action of phosphonoacetaldehyde hydrolase 409–411
LAVELLE, F. *see* MCADAM, M. E. 71–79, 81–87
LINDSAY, J. A. & CREASER, E. H. Purification and properties of histidinol dehydrogenases from psychrophilic, mesophilic and thermophilic bacilli 247–253
LINKER, A. *see* HOVINGH, P. 287–293

MACGIBBON, A. K. H., BUCKLEY, P. D. & BLACKWELL, L. F. Evidence for two-step binding of reduced nicotinamide–adenine dinucleotide to aldehyde dehydrogenase 455–462
MAGEE, A. I., GRANT, D. A. W. & HERMON-TAYLOR, J. The apparent molecular weights of human intestinal aminopeptidase, enterokinase and maltase in native duodenal fluid 583–585
MAJUMDER, G. C. Purification and properties of a nuclear protein kinase from rat mammary gland 469–477
MANDEL, P. *see* HELWIG, J.-J. 127–134
MANJULA, B. N., ACHARYA, A. S. & VITHAYATHIL, P. J. Subtilisin modification of monodeamidated ribonuclease-A 337–345
MANNERVIK, B. *see* ARONSSON, A.-C. 503–509
MANWELL, C. A simplified electrophoretic system for determining molecular weights of proteins 487–495
MARGISON, G. P., MARGISON, J. M. & MONTESANO, R. Accumulation of $O^6$-methylguanine in non-target-tissue deoxyribonucleic acid during chronic administration of dimethylnitrosamine 463–468
MARGISON, J. M. *see* MARGISON, G. P. 463–468
MARINKOVIC, D. V. & MARINKOVIC, J. N. Restoration of $\beta$-galactosidase to *Escherichia coli* M15. Complementation studies 417–423
MARINKOVIC, J. N. *see* MARINKOVIC, D. V. 417–423
MARSH, D. *see* SUTTON, B. J. 177–197; WILLAN, K. J. 199–206
MARTIN, J. B. *see* OGAWA, N. 269–277
MASTERS, C. J. *see* WALSH, T. P. 165–167
MATCHAM, G. W. J. *see* BARTHOLOMEW, B. 575–580
MCADAM, M. E., FOX, R. A., LAVELLE, F. & FIELDEN, E. M. A pulse-radiolysis study of the manganese-containing superoxide dismutase from *Bacillus stearothermophilus*. A kinetic model for the enzyme action 71–79
MCADAM, M. E., LAVELLE, F., FOX, R. A. & FIELDEN, E. M. A pulse-radiolysis study of the manganese-containing superoxide dismutase from *Bacillus stearothermophilus*. Further studies on the properties of the enzyme 81–87
MCLAUGHLIN, A. C. *see* WAIN-HOBSON, S. 227–235
MILLER, A. L., FROST, R. G. & O'BRIEN, J. S. Purified human liver acid $\beta$-D-galactosidases possessing activity towards $G_{M1}$-ganglioside and lactosylceramide 591–594
MOENS, L. & KONDO, M. Characterization of the extracellular haemoglobins of *Artemia salina* 111–119

MONTESANO, R. *see* MARGISON, G. P. 463–468
MULDER, G. J. & SCHOLTENS, E. Phenol sulphotransferase from rat liver *in vivo* and *in vitro*. 2,6-Dichloro-4-nitrophenol as selective inhibitor of sulphation 563–559
MUNDAY, K. A. *see* GILES, I. G. 97–105
MUTO, M. Estimation of gene reiteration from hybridization kinetics in moderate deoxyribonucleic acid excess 19–25

NAKAIE, C. R. *see* NOUAILHETAS, V. L. A. 547–551
NAUDÉ, R. J. & OELOFSEN, W. The isolation and characterization of corticotropin from the pituitary gland of the ostrich *Struthio camelus* 519–523
NICHOLLS, R. G. Purification and steady-state kinetics of adenosine 5′-pyrophosphate sulphurylase from baker's yeast 149–155
NIELSEN, J. *see* HURST, P. L. 33–41
NIMMO, I. A., ATKINS, G. L., STRANGE, R. C. & PERCY-ROBB, I. W. An evaluation of ways of using equilibrium dialysis to quantify the binding of ligand to macromolecule 107–110
NISHIZAWA, Y. *see* KURIHARA, T. 135–140
NOCHUMSON, S., DURBAN, E., KIM, S. & PAIK, W. K. Cytochrome *c*-specific protein methylase III from *Neurospora crassa* 11–18
NOUAILHETAS, V. L. A., NAKAIE, C. R., JULIANO, L. & PAIVA, A. C. M. Effect of lipophilic character on the biological activity of synthetic angiotensin peptides 547–551

O'BRIEN, J. S. *see* MILLER, A. L. 591–594
OELOFSEN, W. *see* NAUDÉ, R. J. 519–523
OFFORD, R. *see* SAUNDERS, D. J. 479–486
OGAWA, N., THOMPSON, T., FRIESEN, H. G., MARTIN, J. B. & BRAZEAU, P. Properties of soluble somatostatin-binding protein 269–277
OLD, J. M. & JONES, D. S. The aminoacylation of transfer ribonucleic acid. Recognition of methionine by *Escherichia coli* methionyl-transfer ribonucleic acid synthetase 367–373
OSAWA, T. *see* YOKOYAMA, K. 431–437

PAIK, W. K. *see* NOCHUMSON, S. 11–18
PAIVA, A. C. M. *see* NOUAILHETAS, V. L. A. 547–551
PARR, D. M. Fragments produced by digestion of human immunoglobulin G subclasses with pepsin in urea 303–308
PAYS, E. Characterization of double-stranded ribonucleic acid sequences present in the initial transcription products of rat liver chromatin 237–245
PECHT, I. *see* WAIN-HOBSON, S. 227–235
PERCY-ROBB, I. W. *see* NIMMO, I. A. 107–110; STRANGE, R. C. 425–429
PERKINS, S. J., DOWER, S. K., GETTINS, P., WAIN-HOBSON, S. & DWEK, R. A. Application of ring-current theory based on the Johnson–Bovey equation to the aromatic amino acids 223–225
PERKINS, S. J. *see also* DOWER, S. K. 207–223
POAT, P. C. *see* GILES, I. G. 97–105
POLYA, G. M. *see* ASHTON, A. R. 27–32
PORTER, R. R. *see* GIGLI, I. 439–446

QUOCK, R. *see* KEENAN, R. W. 405–408

RAGAN, C. I. *see* CROWDER, S. E. 295–301
RICE, N. *see* KEENAN, R. W. 405–408
ROBERTS, D. V. *see* SMITH, G. D. 121–126
RODÉN, L *see* HEANEY-KIERAS, J. 1–9
ROSSIER, J. Acetyl-coenzyme A and coenzyme A analogues. Their effects on rat brain choline acetyltransferase 321–326

SAUNDERS, D. J. & OFFORD, R. Semisynthetic analogues of insulin. The use of *N*-substituted derivatives of methionine as acid-stable protecting groups 479–486
SCHECHTER, I. *see* BURSTEIN, Y. 347–354
SCHOLTENS, E. *see* MULDER, G. J. 553–559
SCHREURS, W. J. *see* SULLIVAN, P. A. 375–383
SHAW, D. J. *see* BARTHOLOMEW, B. 575–580
SHEPHERD, M. G. *see* CHOONG, Y. S. 385–393; HURST, P. L. 33–41; SULLIVAN, P. A. 375–383
SMITH, B. E. *see* CASS, A. E. G. 587–589
SMITH, G. D., ROBERTS, D. V. & DADAY, A. Affinity chromatography and inhibition of chorismate mutase–prephenate dehydrogenase by derivatives of phenylalanine and tyrosine 121–126
SMYTH, D. G. *see* SNELL, C. R. 43–47
SNELL, C. R. & SMYTH, D. G. Biologically active macromolecular forms of oxytocin. [8-Lysine]oxytocin as a suitable ligand 43–47
SPRINGALL, J. P. *see* THOMSON, A. J. 327–336
SRIVASTAVA, S. K. *see* WIKTOROWICZ, J. E. 49–53
STIRLING, D. I. *see* COLBY, J. 395–402
STORER, A. C. & CORNISH-BOWDEN, A. Kinetic evidence for a 'mnemonical' mechanism for rat liver glucokinase 61–69
STORM, E. *see* BJØRKLID, E. 89–96
STRANGE, R. C., CRAMB, R., HAYES, J. D. & PERCY-ROBB, I. W. Partial purification of two lithocholic acid-binding proteins from rat liver 100000*g* supernatants 425–429
STRANGE, R. C. *see also* NIMMO, I. A. 107–110
STRØMME, J. *see* DUGAL, B. 497–502
SULLIVAN, P. A., CHOONG, Y. S., SCHREURS, W. J., CUTFIELD, J. F. & SHEPHERD, M. G. The structure of L-lactate oxidase from *Mycobacterium smegmatis* 375–383
SULLIVAN, P. A. *see also* CHOONG, Y. S. 385–393; HURST, P. L. 33–41
SUNDERLAND, C. A. *see* DOWER, S. K. 207–223; WAIN-HOBSON, S. 227–235; WILLAN, K. J. 199–206
SUTTON, B. J., GETTINS, P., GIVOL, D., MARSH, D., WAIN-HOBSON, S., WILLAN, K. J. & DWEK, R. A. The gross architecture of an antibody-combining site as determined by spin-label mapping 177–197
SUTTON, B. J. *see also* DOWER, S. K. 207–223; WILLAN, K. J. 199–206

TAKAHASHI, Y. *see* KURIHARA, T. 135–140
TANNER, M. J. A., ANSTEE, D. J. & JUDSON, P. A. A carbohydrate-deficient membrane glycoprotein in human erythrocytes of phenotype S–s– 157–161
TERAO, T. *see* YOKOYAMA, K. 431–437
THOMPSON, T. *see* OGAWA, N. 269–277

THOMSON, A. J., BRITTAIN, T., GREENWOOD, C. & SPRINGALL, J. P. Variable-temperature magnetic-circular-dichroism spectra of cytochrome *c* oxidase and its derivatives 327–336
THORNELEY, R. N. F. & CORNISH-BOWDEN, A. Kinetics of nitrogenase of *Klebsiella pneumoniae*. Heterotropic interactions between magnesium–adenosine 5′-diphosphate and magnesium–adenosine 5′-triphosphate 255–262

UELAND, P. M. *see* DØSKELAND, S. O. 561–573

VERONESE, F. M. *see* FONTANA, A. 539–545
VIJAYAKUMAR, E. K. & WEIDEMANN, M. J. Kinetic properties of a magnesium ion- and calcium ion-stimulated adenosine triphosphatase from the outer-membrane fraction of rat spleen mitochondria 355–365
VITA, C. *see* FONTANA, A. 539–545
VITHAYATHIL, P. J. *see* MANJULA, B. N. 337–345
VLIEGENTHART, J. F. G. *see* HALL, R. L. 173–176
VON ZABERN, I. *see* GIGLI, I. 439–446

WADE, R. *see* BRUNDISH, D. E. 169–171
WAIN-HOBSON, S., DOWER, S. K., GETTINS, P., GIVOL, D., MCLAUGHLIN, A. C., PECHT, I., SUNDERLAND, C. A. & DWEK, R. A. Specificity of interactions of hapten side chains with the combining site of the myeloma protein MOPC 315 227–235
WAIN-HOBSON, S. *see also* DOWER, S. K. 207–223; PERKINS, S. J. 223–225; SUTTON, B. J. 177–197; WILLAN, K. J. 199–206
WALEY, S. G. *see* KIENER, P. A. 279–285
WALSH, T. P., CLARKE, F. M. & MASTERS, C. J. Modification of the kinetic parameters of aldolase on binding to the actin-containing filaments of skeletal muscle 165–167
WEIDEMANN, M. J., KOLBUCH-BRADDON, M. E. & HICKMAN, P. E. Effect of substrates and effectors on the reversible inactivation of pig spleen phosphofructokinase by adenosine triphosphate 525–537
WEIDEMANN, M. J. *see also* VIJAYAKUMAR, E. K. 355–365
WHITE, G. F. *see* BARTHOLOMEW, B. 575–580
WIKTOROWICZ, J. E., AWASTHI, Y. C., KUROSKY, A. & SRIVASTAVA, S. K. Purification and properties of human kidney-cortex hexosaminidases A and B 49–53
WILLAN, K. J., MARSH, D., SUNDERLAND, C. A., SUTTON, B. J., WAIN-HOBSON, S., DWEK, R. A. & GIVOL, D. Comparison of the dimensions of the combining sites of the dinitrophenyl-binding immunoglobulin A myeloma proteins MOPC 315, MOPC 460 and XRPC 25 by spin-label mapping 199–206
WILLAN, K. J. *see also* SUTTON, B. J. 177–197
WILSON, M. T. *see* GREENWOOD, C. 413–416
WINTERBOURN, C. C. & CARRELL, R. W. Oxidation of human haemoglobin by copper. Mechanism and suggested role of the thiol group of residue β-93 141–148
WOOD, E. J. *see* HALL, R. L. 173–176
WRIGHT, C. E. *see* DOWER, S. K. 207–223

YOKOYAMA, K., TERAO, T. & OSAWA, T. Isolation and characterization of membrane receptors for pokeweed mitogens from mouse lymphocytes 431–437

# Index of Subjects


Acetyl-coenzyme A, effects of coenzyme A and, and their analogues on the activity of rat brain choline acetyltransferase (Rossier, J.) 321–326

β-D-*N*-Acetylhexosaminidases A and B, kidney-cortex, human, purification and properties of (Wiktorowicz, J. E., Awasthi, Y. C., Kurosky, A. & Srivastava, S. K.) 49–53

Acid β-D-galactosidase, *see* β-D-Galactosidase, acid

Actin, modification of the kinetic parameters of fructose bisphosphate aldolase on binding to ox skeletal-muscle filaments containing (Walsh, T. P., Clarke, F. M. & Masters, C. J.) 165–167

Adenosine, binding of, by ox adrenal-gland-cortex adenosine 3′:5′-cyclic monophosphate-binding proteins (Døskeland, S. O. & Ueland, P. M.) 561–573

Adenosine 3′:5′-cyclic monophosphate-binding proteins, adrenal-gland-cortex, ox, characterization of (Døskeland, S. O. & Ueland, P. M.) 561–573

Adenosine 3′:5′-cyclic monophosphate, isolation and characterization of, from *Kalanchoe daigremontiana* and *Agave americana* (Ashton, A. R. & Polya, G. M.) 27–32

Adenosine diphosphate, effects of the complexes of magnesium ions with adenosine triphosphate and, on the kinetics of the reaction catalysed by *Klebsiella pneumoniae* nitrogenase (Thorneley, R. N. F. & Cornish-Bowden, A.) 255–262

Adenosine diphosphate sulphurylase, *see* Sulphate adenylyltransferase (adenosine diphosphate)

Adenosine triphosphatase, magnesium-and-calcium ion-stimulated, mitochondrial-outer-membrane, spleen, rat, kinetic properties of (Vijayakumar, E. K. & Weidemann, M. J.) 355–365

Adenosine triphosphate, effects of alanine, fructose 1,6-bisphosphate and, on the activity of crab hepatopancreas pyruvate kinase (Giles, I. G., Poat, P. C. & Munday, K. A.) 97–105

Adenosine triphosphate, effects of substrates and effectors on the reversible inactivation by, of pig spleen phosphofructokinase (Weidemann, M. J., Kolbuch-Braddon, M. E. & Hickman, P. E.) 525–537

Adenosine triphosphate, effects of the complexes of magnesium ions with adenosine diphosphate and, on the kinetics of the reaction catalysed by *Klebsiella pneumoniae* nitrogenase (Thorneley, R. N. F. & Cornish-Bowden, A.) 255–262

Adrenal-gland cortex, ox, characterization of adenosine 3′:5′-cyclic monophosphate-binding proteins of (Døskeland, S. O. & Ueland, P. M.) 561–573

Adrenocorticotropin, *see* Corticotropin

*Agave americana*, isolation and characterization of adenosine 3′:5′-cyclic monophosphate from *Kalanchoe daigremontiana* and (Ashton, A. R. & Polya, G. M.) 27–32

Alanine, effects of adenosine triphosphate, fructose 1,6-bisphosphate and, on the activity of crab hepatopancreas pyruvate kinase (Giles, I. G., Poat, P. C. & Munday, K. A.) 97–105

Aldehyde dehydrogenase, liver, sheep, evidence for two-step binding of reduced nicotinamide–adenine dinucleotide to (MacGibbon, A. K. H., Buckley, P. D. & Blackwell, L. F.) 455–462

Aldolase, *see* Fructose bisphosphate aldolase

*n*-Alkanes, ability of *Methylococcus capsulatus* methane mono-oxygenase to catalyse the oxygenation of, and a range of *n*-alkenes, ethers and alicyclic, aromatic and heterocyclic compounds (Colby, J., Stirling, D. I. & Dalton, H.) 395–402

*n*-Alkenes, ability of *Methylococcus capsulatus* methane mono-oxygenase to catalyse the oxygenation of, and a range of *n*-alkanes, ethers and alicyclic, aromatic and heterocyclic compounds (Colby, J., Stirling, D. I. & Dalton, H.) 395–402

Alkylsulphatases, secondary, *Comamonas terrigena* and *Pseudomonas* C12B, inversion of configuration and carbon–oxygen bond cleavage catalysed by (Bartholomew, B., Dodgson, K. S., Matcham, G. W. J., Shaw, D. J. & White, G. F.) 575–580

Amino acid sequence, variability of, of the *N*-terminal extra piece of precursors of immunoglobulin λ-type light chains produced by mouse plasma-cell tumour MOPC 104E, RPC 20 and MOPC 315 cells (Burstein, Y. & Schechter, I.) 347–354

Amino acid sequences, *N*-terminal, determination of, of the α-, β- and γ-chains of component C4 of human complement (Gigli, I., von Zabern, I. & Porter, R. R.) 439–446

Amino acids, aromatic, application of ring-current theory based on the Johnson–Bovey equation to (Perkins, S. J., Dower, S. K., Gettins, P., Wain-Hobson, S. & Dwek, R. A.) 223–225

Aminopeptidase, small-intestinal, human, apparent molecular weights of, and enterokinase and maltase in duodenal fluid (Magee, A. I., Grant, D. A. W. & Hermon-Taylor, J.) 583–585

Angiotensin II, [5-isoleucine]-, effect of lipophilic character on the biological activity of synthetic derivatives of (Nouailhetas, V. L. A., Nakaie, C. R., Juliano, L. & Paiva, A. C. M.) 547–551

*Artemia salina*, *see* Brine shrimp

Arylsulphatases A and B, kidney-cortex, rabbit, purification and properties of (Helwig, J.-J., Farooqui, A. A., Bollack, C. & Mandel, P.) 127–134

Aspartylglucosylaminase, liver, human, purification and properties of (Dugal, B. & Strømme, J.) 497–502

*Aspergillus niger*, purification and properties of a cellulase from (Hurst, P. L., Nielsen, J., Sullivan, P. A. & Shepherd, M. G.) 33–41

*Bacillus caldolyticus*, purification and properties of histidinol dehydrogenases from, and other *Bacillus* spp. (Lindsay, J. A. & Creaser, E. H.) 247–253

*Bacillus cereus*, evidence for the formation of an imine between enzyme and substrate during the reaction catalysed by phosphonoacetaldehyde hydrolase from (La Nauze, J. M., Coggins, J. R. & Dixon, H. B. F.) 409–411

*Bacillus cereus*, study of the conformational motility of β-lactamase I from, by measurement of the exchange of hydrogen atoms with substrates (Kiener, P. A. & Waley, S. G.) 279–285

*Bacillus psychrophilus*, purification and properties of histidinol dehydrogenases from, and other *Bacillus* spp. (Lindsay, J. A. & Creaser, E. H.) 247–253

*Bacillus stearothermophilus*, kinetic model for the reaction catalysed by manganese-containing superoxide dismutase from (McAdam, M. E., Fox, R. A., Lavelle, F. & Fielden, E. M.) 71–79
*Bacillus stearothermophilus*, properties of manganese-containing superoxide dismutase from (McAdam, M. E., Lavelle, F., Fox, R. A. & Fielden, E. M.) 81–87
*Bacillus stearothermophilus*, purification and properties of histidinol dehydrogenases from, and other *Bacillus* spp. (Lindsay, J. A. & Creaser, E. H.) 247–253
*Bacillus subtilis*, purification and properties of histidinol dehydrogenases from, and other *Bacillus* spp. (Lindsay, J. A. & Creaser, E. H.) 247–253
Benzonitrile, purification and properties of *Nocardia* sp. N.C.I.B. 11216 nitrilase that catalyses cleavage of the carbon–nitrogen bond of, and other aromatic nitriles (Harper, D. B.) 309–319
Bile acid-binding proteins, cytosol, liver, rat, two, partial purification of (Strange, R. C., Cramb, R., Hayes, J. D. & Percy-Robb, I. W.) 425–429
Brain, human, purification and properties of the protein component of thromboplastin from (Bjørklid, E. & Storm, E.) 89–96
Brain, ox, use of extraction with non-aqueous chloroform/methanol for the delipidation of, with minimal loss of enzyme activities (Kurihara, T., Nishizawa, Y. & Takahashi, Y.) 135–140
Brain, rat, effects of acetyl-coenzyme A and coenzyme A and their analogues on the activity of choline acetyltransferase from (Rossier, J.) 321–326
Brine shrimp (*Artemia salina*), characterization of the extracellular haemoglobins of (Moens, L. & Kondo, M.) 111–119

Calcium ions, fluorimetric study of the role of, in the stability of thermolysin (Fontana, A., Vita, C., Boccu', E. & Veronese, F. M.) 539–545
Carbohydrate, covalent linkage of protein to, in *Porphyridium cruentum* extracellular proteoglycan (Heaney-Kieras, J., Rodén, L. & Chapman, D. J.) 1–9
Carbohydrate sulphates, specificity of rabbit kidney-cortex arylsulphatases A and B towards (Helwig, J.-J., Farooqui, A. A., Bollack, C. & Mandel, P.) 127–134
*Carcinus maenas*, *see* Crab
Casein kinase, ribosomal, reticulocyte, rabbit, purification and properties of (Issinger, O.-G.) 511–518
Cations, inorganic, effects of amphiphilic cationic drugs and, on the activity of rat liver phosphatidate phosphatase (Bowley, M., Cooling, J., Burditt, S. L. & Brindley, D. N.) 447–454
Cells, myeloma, X5563, mouse, estimation of reiteration of genes in, from kinetics of hybridization of deoxyribonucleic acid and ribonucleic acid in the presence of a moderate excess of deoxyribonucleic acid (Muto, M.) 19–25
Cellulase, *Aspergillus niger*, purification and properties of (Hurst, P. L., Nielsen, J., Sullivan, P. A. & Shepherd, M. G.) 33–41
Choline acetyltransferase, brain, rat, effects of acetyl-coenzyme A and coenzyme A and their analogues on the activity of (Rossier, J.) 321–326
Chondroitin sulphates, specificity of *Flavobacterium heparinum* glycuronidases acting on disaccharides derived from, and other glycosáminoglycans (Hovingh, P. & Linker, A.) 287–293
Chorismate mutase–prephenate dehydrogenase, *Escherichia coli* K12, affinity chromatography of, and inhibition of its activity by derivatives of phenylalanine and tyrosine (Smith, G. D., Roberts, D. V. & Daday, A.) 121–126
Chromatin, liver, rat, characterization of the nucleotide sequences of double-stranded segments of ribonucleic acid present in the initial transcription products of (Pays, E.) 237–245
Chromous ions, reduction by, of mammalian cytochrome *c* oxidase and some of its derivatives (Greenwood, C., Brittain, T., Brunori, M. & Wilson, M. T.) 413–416
Clotting Factor III (thromboplastin), brain, human, purification and properties of (Bjørklid, E. & Storm, E.) 89–96
Coagulation Factor, *see* Clotting Factor
Coenzyme A, effects of acetyl-coenzyme A and, and their analogues on the activity of rat brain choline acetyltransferase (Rossier, J.) 321–326
*Comamonas terrigena*, inversion of configuration and carbon–oxygen bond cleavage catalysed by secondary alkylsulphatases from *Pseudomonas* C12B and (Bartholomew, B., Dodgson, K. S., Matcham, G. W. J., Shaw, D. J. & White, G. F.) 575–580
Complement, human, isolation and structure of component C4 of (Gigli, I., von Zabern, I. & Porter, R. R.) 439–446
Component C4, complement, human, isolation and structure of (Gigli, I., von Zabern, I. & Porter, R. R.) 439–446
Computer program, use of, in an automatic method for the derivation of steady-state rate equations for reactions catalysed by enzymes (Cornish-Bowden, A.) 55–59
Copper ions, mechanism of the oxidation by, of the haem groups of human haemoglobin and suggested role of the thiol group of cysteine residue β-93 (Winterbourn, C. C. & Carrell, R. W.) 141–148
Corticotropin, anterior-pituitary-gland, ostrich, isolation and characterization of (Naudé, R. J. & Oelofsen, W.) 519–523
Corticotropin, human, tritium-labelled, synthesis of, of high specific radioactivity (Brundish, D. E. & Wade, R.) 169–171
Crab (*Carcinus maenas*), interactions of allosteric modifiers with hepatopancreas pyruvate kinase from (Giles, I. G., Poat, P. C. & Munday, K. A.) 97–105
Cyclic adenosine 3′:5′-monophosphate, *see* Adenosine 3′:5′-cyclic monophosphate
2′:3′-Cyclic nucleotide 3′-phosphodiesterase, use of extraction with non-aqueous chloroform/methanol for the delipidation of ox brain with minimal loss of the activities of, and other enzymes (Kurihara, T., Nishizawa, Y. & Takahashi, Y.) 135–140
Cysteine residue β-93, haemoglobin, human, suggested role of the thiol group of, in the oxidation by copper ions of the haem groups of the protein (Winterbourn, C. C. & Carrell, R. W.) 141–148

Cytochrome *c* methylase (protein methylase III), *Neurospora crassa*, partial purification and characterization of (Nochumson, S., Durban, E., Kim, S. & Paik, W. K.) 11–18
Cytochrome *c* oxidase, heart, ox, variable-temperature magnetic-circular-dichroism spectra of, and its derivatives (Thomson, A. J., Brittain, T., Greenwood, C. & Springall, J. P.) 327–336
Cytochrome *c* oxidase, mammalian, reduction by chromous ions of, and some of its derivatives (Greenwood, C., Brittain, T., Brunori, M. & Wilson, M. T.) 413–416
Cytosol, liver, rat, partial purification of two lithocholate-binding proteins from (Strange, R. C., Cramb, R., Hayes, J. D. & Percy-Robb, I. W.) 425–429
Cytosol, properties of somatostatin-binding protein from, of various rat, human and ox tissues (Ogawa, N., Thompson, T., Friesen, H. G., Martin, J. B. & Brazeau, P.) 269–277

Deoxyribonucleic acid, estimation of reiteration of genes in mouse X5563 myeloma cells from kinetics of the hybridization of ribonucleic acid and, in the presence of a moderate excess of deoxyribonucleic acid (Muto, M.) 19–25
Deoxyribonucleic acids, accumulation of $O^6$-methylguanylate residues in, of non-target tissues during chronic administration of dimethylnitrosamine to the rat (Margison, G. P., Margison, J. M. & Montesano, R.) 463–468
2,6-Dichloro-4-nitrophenol, use of, as selective inhibitor of sulphation by phenol sulphotransferase but not of glucuronidation by uridine diphosphate glucuronosyltransferase of rat liver *in vivo* and *in vitro* (Mulder, G. J. & Scholtens, E.) 553–559
Diethyl pyrocarbonate, evidence for an active-site histidine residue in *Mycobacterium smegmatis* L-lactate oxidase obtained by modification of the enzyme with (Choong, Y. S., Shepherd, M. G. & Sullivan, P. A.) 385–393
Dimethylnitrosamine, accumulation of $O^6$-methylguanylate residues in deoxyribonucleic acids of non-target tissues during chronic administration of, to the rat (Margison, G. P., Margison, J. M. & Montesano, R.) 463–468
Dinitrophenyl-binding myeloma protein (immunoglobulin A), geometry of the antibody-combining site of, from mouse plasma-cell tumour MOP 315 cells (Dower, S. K., Wain-Hobson, S., Gettins, P., Givol, D., Jackson, W. R. C., Perkins, S. J., Sunderland, C. A., Sutton, B. J., Wright, C. E. & Dwek, R. A.) 207–223
Dinitrophenyl-binding myeloma protein (immunoglobulin A), gross architecture of the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells determined by spin-label mapping (Sutton, B. J., Gettins, P., Givol, D., Marsh, D., Wain-Hobson, S., Willan, K. J. & Dwek, R. A.) 177–197
Dinitrophenyl-binding myeloma protein (immunoglobulin A), specificity of interactions of hapten side chains with the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells (Wain-Hobson, S., Dower, S. K., Gettins, P., Givol, D., McLaughlin, A. C., Pecht, I., Sunderland, C. A. & Dwek, R. A.) 227–235
Dinitrophenyl-binding myeloma proteins (immunoglobulins A), comparison of the dimensions of the antibody-combining sites of, from mouse plasma-cell tumour MOPC 315, MOPC 460 and XRPC 25 cells determined by spin-label mapping (Willan, K. J., Marsh, D., Sunderland, C. A., Sutton, B. J., Wain-Hobson, S., Dwek, R. A. & Givol, D.) 199–206
Disaccharides, specificity of *Flavobacterium heparinum* glycuronidases acting on, derived from glycosaminoglycans (Hovingh, P. & Linker, A.) 287–293
Dolichol, liver, pig, use of high-performance liquid chromatography for the analysis of, and other polyprenols (Keenan, R. W., Rice, N. & Quock, R.) 405–408
Drugs, cationic, amphiphilic, effects of inorganic cations and, on the activity of rat liver phosphatidate phosphatase (Bowley, M., Cooling, J., Burditt, S. L. & Brindley, D. N.) 447–454
Duodenal fluid, human, apparent molecular weights of small-intestinal aminopeptidase, enterokinase and maltase in (Magee, A. I., Grant, D. A. W. & Hermon-Taylor, J.) 583–585

Enterokinase, small-intestinal, human, apparent molecular weights of, and aminopeptidase and maltase in duodenal fluid (Magee, A. I., Grant, D. A. W. & Hermon-Taylor, J.) 583–585
Enzymes, automatic method for the derivation of steady-state rate equations for reactions catalysed by (Cornish-Bowden, A.) 55–59
Erythrocytes, human, characterization of a carbohydrate-deficient glycoprotein of the membrane of, of phenotype S – s – (Tanner, M. J. A., Anstee, D. J. & Judson, P. A.) 157–161
Erythrocytes, ox, exchange of protons at C-2 of histidine residue 41 of superoxide dismutase from (Cass, A. E. G., Hill, H. A. O., Smith, B. E., Bannister, J. V. & Bannister, W. H.) 587–589
Erythrocytes, pig, purification and properties of glyoxalase I from (Aronsson, A.-C. & Mannervik, B.) 503–509
*Escherichia coli* K12, affinity chromatography of chorismate mutase–prephenate dehydrogenase from, and inhibition of its activity by derivatives of phenylalanine and tyrosine (Smith, G. D., Roberts, D. V. & Daday, A.) 121–126
*Escherichia coli* mutant M15, restoration of β-galactosidase to, by complementation (Marinkovic, D. V. & Marinkovic, J. N.) 417–423
*Escherichia coli*, recognition of methionine by methionyl-transfer ribonucleic acid synthetase from (Old, J. M. & Jones, D. S.) 367–373
Ethoxyformate, *see* Diethyl pyrocarbonate

Factor III, clotting (thromboplastin), brain, human, purification and properties of (Bjørklid, E. & Storm, E.) 89–96
Ficaprenol, *Ficus elastica*, use of high-performance liquid chromatography for the analysis of, and other polyprenols (Keenan, R. W., Rice, N. & Quock, R.) 405–408
*Ficus elastica* (decorative rubber plant), use of high-performance liquid chromatography for the analysis of ficaprenol from, and other polyprenols (Keenan, R. W., Rice, N. & Quock, R.) 405–408

*Flavobacterium heparinum*, specificity of glycuronidases from, acting on disaccharides derived from glycosaminoglycans (Hovingh, P. & Linker, A.) 287–293
Fructose bisphosphate aldolase, skeletal-muscle, ox, modification of the kinetic parameters of, on binding to the actin-containing filaments of skeletal muscle (Walsh, T. P., Clarke, F. M. & Masters, C. J.) 165–167
Fructose 1,6-bisphosphate, effects of alanine, adenosine triphosphate and, on the activity of crab hepatopancreas pyruvate kinase (Giles, I. G., Poat, P. C. & Munday, K. A.) 97–105
Fructose 6-phosphate, effects of, and other substrates and effectors on the inactivation by adenosine triphosphate of pig spleen phosphofructokinase (Weidemann, M. J., Kolbuch-Braddon, M. E. & Hickman, P. E.) 525–537

β-Galactosidase, restoration of, to *Escherichia coli* mutant M15 by complementation (Marinkovic, D. V. & Marinkovic, J. N.) 417–423
β-D-Galactosidases, acid, liver, human, purified, activity of, towards $G_{M1}$-ganglioside and lactosylceramide (Miller, A. L., Frost, R. G. & O'Brien, J. S.) 591–594
Ganglioside, $G_{M1}$-, activity of purified human liver acid β-D-galactosidases towards lactosylceramide and (Miller, A. L., Frost, R. G. & O'Brien, J. S.) 591–594
Genes, estimation of reiteration of, in mouse X5563 myeloma cells, from kinetics of the hybridization of deoxyribonucleic acid and ribonucleic acid in the presence of a moderate excess of deoxyribonucleic acid (Muto, M.) 19–25
Glucokinase, liver, rat, kinetic evidence for a 'mnemonical' mechanism for the reaction catalysed by (Storer, A. C. & Cornish-Bowden, A.) 61–69
Glucuronidation, use of 2,6-dichloro-4-nitrophenol as selective inhibitor of sulphation by phenol sulphotransferase but not of, by uridine diphosphate glucuronosyltransferase of rat liver *in vivo* and *in vitro* (Mulder, G. J. & Scholtens, E.) 553–559
Glutamine residue, *N*-terminal, identification of, before its conversion into 5-oxopyrrolidine-2-carboxylic acid residue in precursors of immunoglobulin λ-type light chains produced by mouse plasma-cell tumour MOPC 104E, RPC 20 and MOPC 315 cells (Burstein, Y. & Schechter, I.) 347–354
Glutathione, reduced, effect of, on the properties of pig erythrocyte glyoxalase I (Aronsson, A.-C. & Mannervik, B.) 503–509
Glutathione *S*-transferase, activity of, exhibited by two partially purified rat liver cytosol lithocholate-binding proteins (Strange, R. C., Cramb, R., Hayes, J. D. & Percy-Robb, I. W.) 425–429
Glycoprotein, carbohydrate-deficient, characterization of, of the membrane of human erythrocytes of phenotype S − s − (Tanner, M. J. A., Anstee, D. J. & Judson, P. A.) 157–161
Glycosaminoglycans, specificity of *Flavobacterium heparinum* glycuronidases on disaccharides derived from (Hovingh, P. & Linker, A.) 287–293
Glycuronidases, *Flavobacterium heparinum*, specificity of, acting on disaccharides derived from glycosaminoglycans (Hovingh, P. & Linker, A.) 287–293
Glyoxalase I, erythrocyte, pig, purification and properties of (Aronsson, A.-C. & Mannervik, B.) 503–509
Haem groups, haemoglobin, human, mechanism of the oxidation by copper ions of, and suggested role of the thiol group of cysteine residue β-93 (Winterbourn, C. C. & Carrell, R. W.) 141–148
Haemocyanins, pulmonate-gastropod, identification of 3-*O*-methylgalactose and 3-*O*-methylmannose in (Hall, R. L., Wood, E. J., Kamerling, J. P., Gerwig, G. J. & Vliegenthart, J. F. G.) 173–176
Haemoglobin, human, mechanism of the oxidation by copper ions of the haem groups of, and suggested role of the thiol group of cysteine residue β-93 (Winterbourn, C. C. & Carrell, R. W.) 141–148
Haemoglobins, extracellular, brine-shrimp, characterization of (Moens, L. & Kondo, M.) 111–119
Haptens, specificity of interactions of side chains of, with the antibody-combining site of dinitrophenyl-binding myeloma protein (immunoglobulin A) from mouse plasma-cell tumour MOPC 315 cells (Wain-Hobson, S., Dower, S. K., Gettins, P., Givol, D., McLaughlin, A. C., Pecht, I., Sunderland, C. A. & Dwek, R. A.) 227–235
Heart, ox, effects of chymotryptic digestion on the structure and catalytic properties of mitochondrial reduced nicotinamide–adenine dinucleotide–ubiquinone oxidoreductase from (Crowder, S. E. & Ragan, C. I.) 295–301
Heart, ox, variable-temperature magnetic-circular-dichroism spectra of cytochrome *c* oxidase from, and its derivatives (Thomson, A. J., Brittain, T., Greenwood, C. & Springall, J. P.) 327–336
*Helix pomatia*, *see* Snail, Roman
Heparan sulphate, specificity of *Flavobacterium heparinum* glycuronidases acting on disaccharides derived from, and other glycosaminoglycans (Hovingh, P. & Linker, A.) 287–293
Heparin, specificity of *Flavobacterium heparinum* glycuronidases acting on disaccharides derived from, and other glycosaminoglycans (Hovingh, P. & Linker, A.) 287–293
Hepatopancreas, crab, interactions of allosteric modifiers with pyruvate kinase from (Giles, I. G., Poat, P. C. & Munday, K. A.) 97–105
Hexosaminidase, *see* β-D-*N*-Acetylhexosaminidase
Histidine residue, active-site, evidence for, in *Mycobacterium smegmatis* L-lactate oxidase obtained by modification of the enzyme with diethyl pyrocarbonate (Choong, Y. S., Shepherd, M. G. & Sullivan, P. A.) 385–393
Histidine residue 41, exchange of protons at C-2 of, of ox erythrocyte superoxide dismutase (Cass, A. E. G., Hill, H. A. O., Smith, B. E., Bannister, J. V. & Bannister, W. H.) 587–589
Histidinol dehydrogenases, purification and properties of, from psychrophilic, mesophilic and thermophilic *Bacillus* spp. (Lindsay, J. A. & Creaser, E. H.) 247–253
Hydrogen ions, exchange of, at C-2 of histine residue 41 of ox erythrocyte superoxide dismutase (Cass, A. E. G., Hill, H. A. O., Smith, B. E., Bannister, J. V. & Bannister, W. H.) 587–589
Hydrogen isotope ($^3H$), synthesis of human corticotropin labelled with, to high specific radioactivity (Brundish, D. E. & Wade, R.) 169–171

Imine, evidence for the formation of, between enzyme and substrate during the reaction catalysed by *Bacillus cereus* phosphonoacetaldehyde hydrolase (La Nauze, J. M., Coggins, J. R. & Dixon, H. B. F.) 409–411
Immunoglobulin A (myeloma protein), dinitrophenyl-binding, geometry of the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells (Dower, S. K., Wain-Hobson, S., Gettins, P., Givol, D., Jackson, W. R. C., Perkins, S. J., Sunderland, C. A., Sutton, B. J., Wright, C. E. & Dwek, R. A.) 207–223
Immunoglobulin A (myeloma protein), dinitrophenyl-binding, gross architecture of the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells determined by spin-label mapping (Sutton, B. J., Gettins, P., Givol, D., Marsh, D., Wain-Hobson, S., Willan, K. J. & Dwek, R. A.) 177–197
Immunoglobulin A (myeloma protein), dinitrophenyl-binding, specificity of interactions of hapten side chains with the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells (Wain-Hobson, S., Dower, S. K., Gettins, P., Givol, D., McLaughlin, A. C., Pecht, I., Sunderland, C. A. & Dwek, R. A.) 227–235
Immunoglobulin G, human fragments produced by peptic digestion of, of different subclasses in the presence of urea (Parr, D. M.) 303–308
Immunoglobulins A (myeloma proteins), dinitrophenyl-binding, comparison of the dimensions of the antibody-combining sites of, from mouse plasma-cell tumour MOPC 315, MOPC 460 and XRPC 25 cells determined by spin-label mapping (Willan, K. J., Marsh, D., Sunderland, C. A., Sutton, B. J., Wain-Hobson, S., Dwek, R. A. & Givol, D.) 199–206
Immunoglobulins, identification of *N*-terminal glutamine residue before its conversion into 5-oxopyrrolidine-2-carboxylic acid residue in precursors of $\lambda$-type light chains of, produced by mouse plasma-cell tumour MOPC 104E, RPC 20 and MOPC 315 cells (Burstein, Y. & Schechter, I.) 347–354
Insulin, use of *N*-substituted derivatives of methionine as acid-stable protecting groups during the semi-synthesis of analogues of (Saunders, D. J. & Offord, R.) 479–486
Intestine, small, human, apparent molecular weights of aminopeptidase, enterokinase and maltase of, in duodenal fluid (Magee, A. I., Grant, D. A. W., & Hermon-Taylor, J.) 583–585
Intestine, small, rat, properties of somatostatin-binding protein from the cytosol fraction of, and other mammalian tissues (Ogawa, N., Thompson, T., Friesen, H. G., Martin, J. B. & Brazeau, P.) 269–277
[5-Isoleucine]angiotensin II, effect of lipophilic character on the biological activity of synthetic derivatives of (Nouailhetas, V. L. A., Nakaie, C. R., Juliano, L. & Paiva, A. C. M.) 547–551

*Kalanchoe daigremontiana*, isolation and characterization of adenosine 3′:5′-cyclic monophosphate from *Agave americana* and (Ashton, A. R. & Polya, G. M.) 27–32
Kidney cortex, human, purification and properties of $\beta$-D-*N*-acetylhexosaminidases A and B from (Wiktorowicz, J. E., Awasthi, Y. C., Kurosky, A. & Srivastava, S. K.) 49–53
Kidney cortex, rabbit, purification and properties of arylsulphatases A and B from (Helwig, J.-J., Farooqui, A. A., Bollack, C. & Mandel, P.) 127–134
Kidney, rat, accumulation of $O^6$-methylguanylate residues in deoxyribonucleic acids of, and other tissues during chronic administration of dimethylnitrosamine (Margison, G. P., Margison, J. M. & Montesano, R.) 463–468
*Klebsiella pneumoniae*, effects of magnesium–adenosine diphosphate and magnesium–adenosine triphosphate complex ions on the kinetics of the reaction catalysed by nitrogenase from (Thorneley, R. N. F. & Cornish-Bowden, A.) 255–262

$\beta$-Lactamase I, *Bacillus cereus*, study of the conformational motility of, by measurement of the exchange of hydrogen atoms with substrates (Kiener, P. A. & Waley, S. G.) 279–285
Lactate dehydrogenase, sheep, simplified electrophoretic system for determination of the molecular weights of isoenzymes of, and other proteins (Manwell, C.) 487–495
L-Lactate oxidase, *Mycobacterium smegmatis*, evidence for an active-site histidine residue in, obtained by modification of the enzyme with diethyl pyrocarbonate (Choong, Y. S., Shepherd, M. G. & Sullivan, P. A.) 385–393
L-Lactate oxidase, *Mycobacterium smegmatis*, structure of (Sullivan, P. A., Choong, Y. S., Schreurs, W. J., Cutfield, J. F. & Shepherd, M. G.) 375–383
Lactosylceramide, activity of purified human liver acid $\beta$-D-galactosidases towards $G_{M1}$-ganglioside and (Miller, A. L., Frost, R. G. & O'Brien, J. S.) 591–594
Ligand, evaluation of ways of using equilibrium dialysis to quantify the binding of, to macromolecule (Nimmo, I. A., Atkins, G. L., Strange, R. C. & Percy-Robb, I. W.) 107–110
Light chains, $\lambda$-type, immunoglobulin, identification of *N*-terminal glutamine residue before its conversion into 5-oxopyrrolidine-2-carboxylic acid residue in precursors of, produced by mouse plasma-cell tumour MOPC 104E, RPC 20 and MOPC 315 cells (Burstein, Y. & Schechter, I.) 347–354
Lipids, use of extraction with non-aqueous chloroform/methanol for the removal of, from ox brain with minimal loss of enzyme activities (Kurihara, T., Nishizawa, Y. & Takahashi, Y.) 135–140
Lithocholate-binding proteins, cytosol, liver, rat, two, partial purification of (Strange, R. C., Cramb, R., Hayes, J. D. & Percy-Robb, I. W.) 425–429
Liver, human, activity towards $G_{M1}$-ganglioside and lactosylceramide of acid $\beta$-D-galactosidases purified from (Miller, A. L., Frost, R. G. & O'Brien, J. S.) 591–594
Liver, human, purification and properties of aspartylglucosylaminase from (Dugal, B. & Strømme, J.) 497–502
Liver, pig, use of high-performance liquid chromatography for the analysis of dolichol from, and other polyprenols (Keenan, R. W., Rice, N. & Quock, R.) 405–408

Liver, rat, accumulation of $O^6$-methylguanylate residues in deoxyribonucleic acids of, and other tissues during chronic administration of dimethylnitrosamine (Margison, G. P., Margison, J. M. & Montesano, R.) 463–468

Liver, rat and human, properties of somatostatin-binding protein from the cytosol fraction of, and other mammalian tissues (Ogawa, N., Thompson, T., Friesen, H. G., Martin, J. B. & Brazeau, P.) 269–277

Liver, rat, characterization of the nucleotide sequences of double-stranded segments of ribonucleic acid present in the initial transcription products of chromatin from (Pays, E.) 237–245

Liver, rat, effects of amphiphilic cationic drugs and inorganic cations on the activity of phosphatidate phosphatase from (Bowley, M., Cooling, J., Burditt, S. L. & Brindley, D. N.) 447–454

Liver, rat, kinetic evidence for a 'mnemonical' mechanism for the reaction catalysed by glucokinase from (Storer, A. C. & Cornish-Bowden, A.) 61–69

Liver, rat, partial purification of two lithocholate-binding proteins from cytosol of (Strange, R. C., Cramb, R., Hayes, J. D. & Percy-Robb, I. W.) 425–429

Liver, rat, use of 2,6-dichloro-4-nitrophenol as selective inhibitor of sulphation by phenol sulphotransferase but not of glucuronidation by uridine diphosphate glucuronosyltransferase of, *in vivo* and *in vitro* (Mulder, G. J. & Scholtens, E.) 553–559

Liver, sheep, evidence for two-step binding of reduced nicotinamide–adenine dinucleotide to aldehyde dehydrogenase from (MacGibbon, A. K. H., Buckley, P. D. & Blackwell, L. F.) 455–462

Lung, rat, accumulation of $O^6$-methylguanylate residues in deoxyribonucleic acids of, and other tissues during chronic administration of dimethylnitrosamine (Margison, G. P., Margison, J. M. & Montesano, R.) 463–468

*Lymnaea stagnalis*, *see* Snail, freshwater

Lymphocytes, mouse, isolation and characterization of plasma-membrane pokeweed-mitogen-receptor proteins from (Yokoyama, K., Terao, T. & Osawa, T.) 431–437

[8-Lysine]oxytocin, suitability of, as a ligand for the preparation of immobilized forms of oxytocin (Snell, C. R. & Smyth, D. G.) 43–47

Macromolecule, evaluation of ways of using equilibrium dialysis to quantify the binding of ligand to (Nimmo, I. A., Atkins, G. L., Strange, R. C. & Percy-Robb, I. W.) 107–110

Magnesium ions, effects of the complexes of, with adenosine diphosphate and adenosine triphosphate on the kinetics of the reaction catalysed by *Klebsiella pneumoniae* nitrogenase (Thorneley, R. N. F. & Cornish-Bowden, A.) 255–262

Maltase, small-intestinal, human, apparent molecular weights of, and aminopeptidase and enterokinase in duodenal fluid (Magee, A. I., Grant, D. A. W. & Hermon-Taylor, J.) 583–585

Mammary gland, rat, purification and properties of a nuclear protein kinase from (Majumder, G. C.) 469–477

Manganese, kinetic model for the reaction catalysed by *Bacillus stearothermophilus* superoxide dismutase containing (McAdam, M. E., Fox, R. A., Lavelle, F. & Fielden, E. M.) 71–79

Manganese, properties of *Bacillus stearothermophilus* superoxide dismutase containing (McAdam, M. E. Lavelle, F., Fox, R. A. & Fielden, E. M.) 81–87

Membrane, erythrocyte, human characterization of a carbohydrate-deficient glycoprotein of, of phenotype S−s− (Tanner, M. J. A., Anstee, D. J. & Judson, P. A.) 157–161

Membrane, lymphocyte, mouse, isolation and characterization of pokeweed-mitogen-receptor proteins from (Yokoyama, K., Terao, T. & Osawa, T.) 431–437

Membrane, outer, mitochondrial, spleen, rat, kinetic properties of magnesium-and-calcium ion-stimulated adenosine triphosphatase from (Vijayakumar, E. K. & Weidemann, M. J.) 355–365

Methane mono-oxygenase, *Methylococcus capsulatus*, ability of, to catalyse the oxygenation of a range of *n*-alkanes, *n*-alkenes, ethers and alicyclic, aromatic and heterocyclic compounds (Colby, J., Stirling, D. I. & Dalton, H.) 395–402

Methanol, ability of *Methylococcus capsulatus* methane mono-oxygenase to catalyse the oxygenation of, and a range of *n*-alkanes, *n*-alkenes, ethers and alicyclic, aromatic and heterocyclic compounds (Colby, J., Stirling, D. I. & Dalton, H.) 395–402

Methionine, recognition of, by *Escherichia coli* methionyl-transfer ribonucleic acid synthetase (Old, J. M. & Jones, D. S.) 367–373

Methionine, use of *N*-substituted derivatives of, as acid-stable protecting groups during the semi-synthesis of analogues of insulin (Saunders, D. J. & Offord, R.) 479–486

Methionyl-transfer ribonucleic acid synthetase, *Escherichia coli*, recognition of methionine by (Old, J. M. & Jones, D. S.) 367–373

3-*O*-Methylgalactose, identification of 3-*O*-methylmannose and, in pulmonate-gastropod haemocyanins (Hall, R. L., Wood, E. J., Kamerling, J. P., Gerwig, G. J. & Vliegenthart, J. F. G.) 173–176

$O^6$-Methylguanylate residues, accumulation of, in deoxyribonucleic acids of non-target tissues during chronic administration of dimethylnitrosamine to the rat (Margison, G. P., Margison, J. M. & Montesano, R.) 463–468

3-*O*-Methylmannose, identification of 3-*O*-methylgalactose and, in pulmonate-gastropod haemocyanins (Hall, R. L., Wood, E. J., Kamerling, J. P., Gerwig, G. J. & Vliegenthart, J. F. G.) 173–176

*Methylococcus capsulatus*, ability of methane mono-oxygenase to catalyse the oxygenation of a range of *n*-alkanes, *n*-alkenes, ethers and alicyclic, aromatic and heterocyclic compounds (Colby, J., Stirling, D. I. & Dalton, H.) 395–402

Microsomal fraction, brain, human, purification and properties of the protein component of thromboplastin from (Bjørklid, E. & Storm, E.) 89–96

Mitochondria, heart, ox, effects of chymotryptic digestion on the structure and catalytic properties of reduced nicotinamide–adenine dinucleotide–ubiquinone oxidoreductase from (Crowder, S. E. & Ragan, C. I.) 295–301

Mitochondria, spleen, rat, kinetic properties of magnesium-and-calcium ion-stimulated adenosine triphosphatase from the outer-membrane fraction of (Vijayakumar, E. K. & Weidemann, M. J.) 355–365
Mitogen (pokeweed)-receptor proteins, membrane, lymphocyte, mouse, isolation and characterization of (Yokoyama, K., Terao, T. & Osawa, T.) 431–437
Muscle, skeletal, ox, modification of the kinetic parameters of fructose bisphosphate aldolase on binding to the actin-containing filaments of (Walsh, T. P., Clarke, F. M. & Masters, C. J.) 165–167
Muscle, skeletal, rat, properties of somatostatin-binding protein from the cytosol fraction of, and other mammalian tissues (Ogawa, N., Thompson, T., Friesen, H. G., Martin, J. B. & Brazeau, P.) 269–277
*Mycobacterium smegmatis*, evidence for an active-site histidine residue in L-lactate oxidase from, obtained by modification of the enzyme with diethyl pyrocarbonate (Choong, Y. S., Shepherd, M. G. & Sullivan, P. A.) 385–393
*Mycobacterium smegmatis*, structure of L-lactate oxidase from (Sullivan, P. A., Choong, Y. S., Schreurs, W. J., Cutfield, J. F. & Shepherd, M. G.) 375–383
Myeloma cells, X5563, mouse, estimation of reiteration of genes in, from kinetics of hybridization of deoxyribonucleic acid and ribonucleic acid in the presence of a moderate excess of deoxyribonucleic acid (Muto, M.) 19–25
Myeloma protein (immunoglobulin A), dinitrophenyl-binding, geometry of the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells (Dower, S. K., Wain-Hobson, S., Gettins, P., Givol, D., Jackson, W. R. C., Perkins, S. J., Sunderland, C. A., Sutton, B. J., Wright, C. E. & Dwek, R. A.) 207–223
Myeloma protein (immunoglobulin A), dinitrophenyl-binding, gross architecture of the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells determined by spin-label mapping (Sutton, B. J., Gettins, P., Givol, D., Marsh, D., Wain-Hobson, S., Willan, K. J. & Dwek, R. A.) 177–197
Myeloma protein (immunoglobulin A), dinitrophenyl-binding, specificity of interactions of hapten side chains with the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells (Wain-Hobson, S., Dower, S. K., Gettins, P., Givol, D., McLaughlin, A. C., Pecht, I., Sunderland, C. A. & Dwek, R. A.) 227–235
Myeloma proteins, human, fragments produced by peptic digestion of, of different immunoglobulin G subclasses in the presence of urea (Parr, D. M.) 303–308
Myeloma proteins, identification of *N*-terminal glutamine residue before its conversion into 5-oxopyrrolidine-2-carboxylic acid residue in precursors of λ-type light chains of, produced by mouse plasma-cell tumour MOPC 104E, RPC 20 and MOPC 315 cells (Burstein, Y. & Schechter, I.) 347–354
Myeloma proteins (immunoglobulins A), dinitrophenyl-binding, comparison of the dimensions of the antibody-combining sites of, from mouse plasma-cell tumour MOPC 315, MOPC 460 and XRPC 25 cells determined by spin-label mapping (Willan, K. J., Marsh, D., Sunderland, C. A., Sutton, B. J., Wain-Hobson, S., Dwek, R. A. & Givol, D.) 199–206
*Neurospora crassa*, partial purification and characterization of cytochrome *c*-specific protein methylase III from (Nochumson, S., Durban, E., Kim, S. & Paik, W. K.) 11–18
Nicotinamide–adenine dinucleotide, reduced, evidence for two-step binding of, to sheep liver aldehyde dehydrogenase (MacGibbon, A. K. H., Buckley, P. D. & Blackwell, L. F.) 455–462
Nicotinamide–adenine dinucleotide (reduced)–ubiquinone oxidoreductase, mitochondrial, heart, ox, effects of chymotryptic digestion on the structure and catalytic properties of (Crowder, S. E. & Ragan, C. I.) 295–301
Nitrilase, *Nocardia* sp. N.C.I.B. 11216, purification and properties of, that catalyses cleavage of the carbon–nitrogen bond of aromatic nitriles (Harper, D. B.) 309–319
Nitriles, aromatic, purification and properties of *Nocardia* sp. N.C.I.B. 11216 nitrilase that catalyses cleavage of the carbon–nitrogen bond of (Harper, D. B.) 309–319
Nitrogenase, *Klebsiella pneumoniae*, effects of magnesium–adenosine diphosphate and magnesium–adenosine triphosphate complex ions on the kinetics of the reaction catalysed by (Thorneley, R. N. F. & Cornish-Bowden, A.) 255–262
Nitrosobenzene, formation of *N*-phenylglycolhydroxamate from, catalysed by yeast transketolase (Corbett, M. D. & Chipko, B. R.) 263–267
*Nocardia* sp. N.C.I.B. 11216, purification and properties of nitrilase from, that catalyses cleavage of the carbon–nitrogen bond of aromatic nitriles (Harper, D. B.) 309–319
Nuclei, mammary-gland, rat, purification and properties of a protein kinase from (Majumder, G. C.) 469–477

Ostrich (*Struthio camelus*), isolation and characterization of corticotropin from the anterior pituitary gland of (Naudé, R. J. & Oelofsen, W.) 519–523
5-Oxopyrrolidine-2-carboxylic acid residue, identification of *N*-terminal glutamine residue before its conversion into, in precursors of immunoglobulin λ-type light chains produced by mouse plasma-cell tumour MOPC 104E, RPC 20 and MOPC 315 cells (Burstein, Y. & Schechter, I.) 347–354
Oxytocin-binding proteins, suitability of [8-lysine]-oxytocin as a ligand for the preparation of immobilized forms of oxytocin for affinity chromatography (Snell, C. R. & Smyth, D. G.) 43–47
Oxytocin, [8-lysine]-, suitability of, as a ligand for the preparation of immobilized forms of oxytocin (Snell, C. R. & Smyth, D. G.) 43–47

Pancreas, ox, modification by limited subtilisin digestion of the monodeamidated derivative of ribonuclease A from (Manjula, B. N., Acharya, A. S. & Vithayathil, P. J.) 337–345
Penicillinase (β-lactamase I), *Bacillus cereus*, study of the conformational motility of, by measurement of the exchange of hydrogen atoms with substrates (Kiener, P. A. & Waley, S. G.) 279–285
Penicillins, study of the conformational motility of *Bacillus cereus* β-lactamase I by measurement of the exchange of hydrogen atoms with, as substrates (Kiener, P. A. & Waley, S. G.) 279–285

Phenol sulphotransferase, use of 2,6-dichloro-4-nitrophenol as selective inhibitor of sulphation by, but not of glucuronidation by uridine diphosphate glucuronosyltransferase of rat liver *in vivo* and *in vitro* (Mulder, G. J. & Scholtens, E.) 553–559

Phenylalanine, inhibition by derivatives of tyrosine and, of the activity of *Escherichia coli* K12 chorismate mutase–prephenate dehydrogenase (Smith, G. D., Roberts, D. V. & Daday, A.) 121–126

*N*-Phenylglycolhydroxamate, formation of, from nitrosobenzene catalysed by yeast transketolase (Corbett, M. D. & Chipko, B. R.) 263–267

Phosphatidate phosphatase, liver, rat, effects of amphiphilic cationic drugs and inorganic cations on the activity of (Bowley, M., Cooling, J., Burditt, S. L. & Brindley, D. N.) 447–454

Phosphofructokinase, spleen, pig, effects of substrates and effectors on the reversible inactivation by adenosine triphosphate of (Weidemann, M. J., Kolbuch-Braddon, M. E. & Hickman, P. E.) 525–537

Phosphonoacetaldehyde hydrolase, *Bacillus cereus*, evidence for the formation of an imine between enzyme and substrate during the reaction catalysed by (La Nauze, J. M., Coggins, J. R. & Dixon, H. B. F.) 409–411

Pituitary gland, anterior, ostrich, isolation and characterization of corticotropin from (Naudé, R. J. & Oelofsen, W.) 519–523

Pituitary gland, anterior, rat, human and ox, properties of somatostatin-binding protein from the cytosol fraction of, and other mammalian tissues (Ogawa, N., Thompson, T., Friesen, H. G., Martin, J. B. & Brazeau, P.) 269–277

Plasma-cell tumour MOPC 104E, RPC 20 and MOPC 315 cells, mouse, identification of *N*-terminal glutamine residue before its conversion into 5-oxopyrrolidine-2-carboxylic acid residue in precursors of immunoglobulin $\lambda$-type light chains produced by (Burstein, Y. & Schechter, I.) 347–354

Plasma-cell tumour MOPC 315 cells, mouse, geometry of the antibody-combining site of dinitrophenyl-binding myeloma protein (immunoglobulin A) from (Dower, S. K., Wain-Hobson, S., Gettins, P., Givol, D., Jackson, W. R. C., Perkins, S. J., Sunderland, C. A., Sutton, B. J., Wright, C. E. & Dwek, R. A.) 207–223

Plasma-cell tumour MOPC 315 cells, mouse, gross architecture of the antibody-combining site of dinitrophenyl-binding myeloma protein (immunoglobulin A) from, determined by spin-label mapping (Sutton, B. J., Gettins, P., Givol, D., Marsh, D., Wain-Hobson, S., Willan, K. J. & Dwek, R. A. 177–197

Plasma-cell tumour MOPC 315 cells, specificity of interactions of hapten side chains with the antibody-combining site of dinitrophenyl-binding myeloma protein (immunoglobulin A) from (Wain-Hobson, S., Dower, S. K., Gettins, P., Givol, D., McLaughlin, A. C., Pecht, I., Sunderland, C. A. & Dwek, R. A.) 227–235

Plasma-cell tumour MOPC 315, MOPC 460 and XRPC 25, cells, mouse, comparison of the dimensions of the antibody-combining sites of dinitrophenyl-binding myeloma proteins (immunoglobulins A) from, determined by spin-label mapping (Willan, K. J., Marsh, D., Sunderland, C. A., Sutton, B. J., Wain-Hobson, S., Dwek, R. A. & Givol, D.) 199–206

Pokeweed-mitogen-receptor proteins, membrane, lymphocyte, mouse, isolation and characterization of (Yokoyama, K., Terao, T. & Osawa, T.) 431–437

Polyprenols, use of high-performance liquid chromatography for the analysis of (Keenan, R. W., Rice, N. & Quock, R.) 405–408

*Porphyridium cruentum*, covalent linkage of protein to carbohydrate in the extracellular proteoglycan from (Heaney-Kieras, J., Rodén, L. & Chapman, D. J.) 1–9

Prephenate dehydrogenase, *Escherichia coli* K12, affinity chromatography of, and inhibition of its activity by derivatives of phenylalanine and tyrosine (Smith, G. D., Roberts, D. V. & Daday, A.) 121–126

Protein component, purification and properties of, of human brain thromboplastin (Bjørklid, E. & Storm, E.) 89–96

Protein, covalent linkage of carbohydrate to, in *Porphyridium cruentum* extracellular proteoglycan (Heaney-Kieras, J., Rodén, L. & Chapman, D. J.) 1–9

Protein kinase, nuclear, mammary-gland, rat, purification and properties of (Majumder, G. C.) 469–477

Protein kinase, ribosomal, reticulocyte, rabbit, purification and properties of (Issinger, O.-G.) 511–518

Protein kinases, adenosine 3′:5′-cyclic monophosphate-dependent, adrenal-gland-cortex, ox, characterization of, and other adenosine 3′:5′-cyclic monophosphate-binding proteins (Døskerland, S. O. & Ueland, P. M.) 561–573

Protein methylase III, cytochrome *c*-specific, *Neurospora crassa*, partial purification and characterization of (Nochumson, S., Durban, E., Kim, S. & Paik, W. K.) 11–18

Protein, myeloma (immunoglobulin A), dinitrophenyl-binding, geometry of the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells (Dower, S. K., Wain-Hobson, S., Gettins, P., Givol, D., Jackson, W. R. C., Perkins, S. J., Sunderland, C. A., Sutton, B. J., Wright, C. E. & Dwek, R. A.) 207–223

Protein, myeloma (immunoglobulin A), dinitrophenyl-binding, gross architecture of the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells determined by spin-label mapping (Sutton, B. J., Gettins, P., Givol, D., Marsh, D., Wain-Hobson, S., Willan, K. J. & Dwek, R. A.) 177–197

Protein, myeloma (immunoglobulin A), dinitrophenyl-binding, specificity of interactions of hapten side chains with the antibody-combining site of, from mouse plasma-cell tumour MOPC 315 cells (Wain-Hobson, S., Dower, S. K., Gettins, P., Givol, D., McLaughlin, A. C., Pecht, I., Sunderland, C. A. & Dwek, R. A.) 227–235

Protein–polysaccharide, *see* Proteoglycan

Protein, somatostatin-binding, properties of, from the cytosol fraction of various rat, human and ox tissues (Ogawa, N., Thompson, T., Friesen, H. G., Martin, J. B. & Brazeau, P.) 269–277

Proteins, adenosine 3′:5′-cyclic monophosphate-binding, adrenal-gland-cortex, ox, characterization of (Døskerland, S. O. & Ueland, P. M.) 561–573

Proteins, lithocholate-binding, cytosol, liver, rat, two, partial purification of (Strange, R. C., Cramb, R., Hayes, J. D. & Percy-Robb, I. W.) 425–429

Proteins, myeloma, human, fragments produced by peptic digestion of, of different immunoglobulin G subclasses in the presence of urea (Parr, D. M.) 303–308

Proteins, myeloma, identification of *N*-terminal glutamine residue before its conversion into 5-oxopyrrolidine-2-carboxylic acid residue in precursors of λ-type light chains of, produced by mouse plasma-cell tumour MOPC 104E, RPC 20 and MOPC 315 cells (Burstein, Y. & Schechter, I.) 347–354

Proteins, myeloma (immunoglobulins A), dinitrophenyl-binding, comparison of the dimensions of the antibody-combining sites of, from mouse plasma-cell tumour MOPC 315, MOPC 460 and XRPC 25 cells determined by spin-label mapping (Willan, K. J., Marsh, D., Sunderland, C. A., Sutton, B. J., Wain-Hobson, S., Dwek, R. A. & Givol, D.) 199–206

Proteins, oxytocin-binding, suitability of [8-lysine]oxytocin as a ligand for the preparation of immobilized forms of oxytocin for affinity chromatography (Snell, C. R. & Smyth, D. G.) 43–47

Proteins, pokeweed-mitogen-receptor, membrane, lymphocyte, mouse isolation and characterization of (Yokoyama, K., Terao, T. & Osawa, T.) 431–437

Proteins, simplified electrophoretic system for determination of the molecular weights of (Manwell, C.) 487–495

Proteoglycan, extracellular, *Porphyridium cruentum*, covalent linkage of protein to carbohydrate in (Heaney-Kieras, J., Rodén, L. & Chapman, D. J.) 1–9

Protons, exchange of, at C-2 of histidine residue 41 of ox erythrocyte superoxide dismutase (Cass, A. E. G., Hill, H. A. O., Smith, B. E., Bannister, J. V. & Bannister, W. H.) 587–589

*Pseudomonas* C12B, inversion of configuration and carbon–oxygen bond cleavage catalysed by secondary alkylsulphatases from *Comamonas terrigena* and (Bartholomew, B., Dodgson, K. S., Matcham, G. W. J., Shaw, D. J. & White, G. F.) 575–580

Pyroglutamic acid, *see* 5-Oxopyrrolidine-2-carboxylic acid

Pyrrolid-2-one-5-carboxylic acid, *see* 5-Oxopyrrolidine-2-carboxylic acid

Pyruvate kinase, hepatopancreas, crab, interactions of allosteric modifiers with (Giles, I. G., Poat, P. C. & Munday, K. A.) 97–105

Red blood cells, *see* Erythrocytes

Reticulocytes, rabbit, purification and properties of ribosomal casein kinase from (Issinger, O.-G.) 511–518

Ribonuclease A, pancreatic, ox, modification by limited subtilisin digestion of the monodeamidated derivative of (Manjula, B. N., Acharya, A. S. & Vithayathil, P. J.) 337–345

Ribonucleic acid, characterization of the nucleotide sequences of double-stranded segments of, present in the initial transcription products of rat liver chromatin (Pays, E.) 237–245

Ribonucleic acid, estimation of reiteration of genes in mouse X5563 myeloma cells from kinetics of the hybridization of deoxyribonucleic acid and, in the presence of a moderate excess of deoxyribonucleic acid (Muto, M.) 19–25

Ribonucleic acid polymerase B, liver, rat, characterization of the nucleotide sequences of double-stranded segments of ribonucleic acid present in the initial transcription products formed by, in the presence of rat liver chromatin (Pays, E.) 237–245

Ribosomes, reticulocyte, rabbit, purification and properties of casein kinase from (Issinger, O.-G.) 511–518

*Saccharomyces cerevisiae*, purification of sulphate adenylyltransferase (adenosine diphosphate) from, and steady-state kinetics of the reaction catalysed by the enzyme (Nicholls, R. G.) 149–155

Sandhoff's disease, purification and properties of human kidney-cortex β-D-*N*-acetylhexosaminidases A and B for possible use in the treatment of (Wiktorowicz, J. E., Awasthi, Y. C., Kurosky, A. & Srivastava, S. K.) 49–53

Skeletal muscle, *see* Muscle, skeletal

Small intestine, *see* Intestine, small

Snail, freshwater (*Lymnaea stagnalis*), identification of 3-*O*-methylgalactose and 3-*O*-methylmannose in haemocyanins of the Roman snail (*Helix pomatia*) and (Hall, R. L., Wood, E. J., Kamerling, J. P., Gerwig, G. J. & Vliegenthart, J. F. G.) 173–176

Snail, Roman (*Helix pomatia*), identification of 3-*O*-methylgalactose and 3-*O*-methylmannose in haemocyanins of a freshwater snail (*Lymnaea stagnalis*) and (Hall, R. L., Wood, E. J., Kamerling, J. P., Gerwig, G. J. & Vliegenthart, J. F. G.) 173–176

Somatostatin-binding protein, properties of, from the cytosol fraction of various rat, human and ox tissues (Ogawa, N., Thompson, T., Friesen, H. G., Martin, J. B. & Brazeau, P.) 269–277

Spleen, pig, effects of substrates and effectors on the reversible inactivation by adenosine triphosphate of phosphofructokinase from (Weidemann, M. J., Kolbuch-Braddon, M. E. & Hickman, P. E.) 525–537

Spleen, rat, kinetic properties of magnesium-and-calcium ion-stimulated adenosine triphosphatase from the outer-membrane fraction of mitochondria from (Vijayakumar, E. K. & Weidemann, M. J.) 355–365

*Struthio camelus*, *see* Ostrich

Sulphate adenylyltransferase (adenosine diphosphate), baker's-yeast, purification of, and steady-state kinetics of the reaction catalysed by the enzyme (Nicholls, R. G.) 149–155

Sulphation, use of 2,6-dichloro-4-nitrophenol as selective inhibitor of, by phenol sulphotransferase but not of glucuronidation by uridine diphosphate glucuronosyltransferase of rat liver *in vivo* and *in vitro* (Mulder, G. J. & Scholtens, E.) 553–559

Supernatant fraction, *see* Cytosol

Superoxide dismutase, erythrocyte, ox, exchange of protons at C-2 of histidine residue 41 of (Cass, A. E. G., Hill, H. A. O., Smith, B. E., Bannister, J. V. & Bannister, W. H.) 587–589

Superoxide dismutase, manganese-containing, *Bacillus stearothermophilus*, kinetic model of the reaction catalysed by (McAdam, M. E., Fox, R. A., Lavelle, F. & Fielden, E. M.) 71–79
Superoxide dismutase, manganese-containing, *Bacillus stearothermophilus*, properties of (McAdam, M. E., Lavelle, F., Fox, R. A. & Fielden, E. M.) 81–87

Tay–Sachs disease, purification and properties of human kidney-cortex β-D-*N*-acetylhexosaminidase A for possible use in the treatment of (Wiktorowicz, J. E., Awasthi, Y. C., Kurosky, A. & Srivastava, S. K.) 49–53
Thermolysin, fluorimetric study of the role of calcium ions in the stability of (Fontana, A., Vita, C., Boccu', E. & Veronese, F. M.) 539–545
Thiol group, suggested role of, of cysteine residue β-93 in the oxidation by copper ions of the haem groups of human haemoglobin (Winterbourn, C. C. & Carrell, R. W.) 141–148
Thromboplastin (clotting Factor III), brain, human, purification and properties of (Bjørklid, E. & Storm, E.) 89–96
Transketolase, yeast, formation of *N*-phenylglycolhydroxamate from nitrosobenzene catalysed by (Corbett, M. D. & Chipko, B. R.) 263–267
Tritium isotope ($^3$H), synthesis of human corticotropin labelled with, to high specific radioactivity (Brundish, D. E. & Wade, R.) 169–171
Tyrosine, inhibition by derivatives of phenylalanine and, of the activity of *Escherichia coli* K12 chorismate mutase–prephenate dehydrogenase (Smith, G. D., Roberts, D. V. & Daday, A.) 121–126

Ubiquinone reductase, nicotinamide–adenine dinucleotide-linked, mitochondrial, heart, ox, effects of chymotryptic digestion on the structure and catalytic properties of (Crowder, S. E. & Ragan, C. I.) 295–301
Uridine diphosphate glucuronosyltransferase, use of 2,6-dichloro-4-nitrophenol as selective inhibitor of sulphation by phenol sulphotransferase but not of glucuronidation by, of rat liver *in vivo* and *in vitro* (Mulder, G. J. & Scholtens, E.) 553–559

Yeast, baker's (*Saccharomyces cerevisiae*), purification of sulphate adenylyltransferase (adenosine diphosphate) from, and steady-state kinetics of the reaction catalysed by the enzyme (Nicholls, R. G.) 149–155
Yeast, formation of *N*-phenylglycolhydroxamate from nitrosobenzene catalysed by transketolase from (Corbett, M. D. & Chipko, B. R.) 263–267

BJECTS

icotropin
Brundish,

hine and,
orismate
G. D.,

dinucleo-
f chymo-
c proper-
301
use of
bitor of
t not of
*in vitro*

cation of
nosphate)
catalysed

ate from
om (Cor-

1977